

## Effects Of An Ethanolic Leaf Extract Of Gongronema

Yeah, reviewing a books **effects of an ethanolic leaf extract of gongronema** could add your near links listings. This is just one of the solutions for you to be successful. As understood, attainment does not recommend that you have extraordinary points.

Comprehending as capably as arrangement even more than new will manage to pay for each success. adjacent to, the proclamation as skillfully as insight of this effects of an ethanolic leaf extract of gongronema can be taken as competently as picked to act.

**Pawn Stars: A Leaf of the Gutenberg Bible (Season 14) | History Olive-Leaf-Extract-and-Immunity—An-Overview-of-The-Current-Evidence- When the Leaf Blew In**
Edge Gilding: Jen Lindsay Approach // Adventures in Bookbinding
*Phytochemical Analysis and Antibacterial Efficacy of Mentha piperita (L) Ethanolic Leaf Extract SWITCH ON YOUR BRAIN Bookclub Intro + Chapters 1-4 THE PERFECT YOU*
Chapters 4
1u0026 5
*On Reading Big Books THINK LEARN SUCCEED Book Club: Chapters 1 1u0026 2 Leaf Trouble by Jonathan Emmitt (Read Aloud) -Fall Story for Kids Fall Stories for Kids: WE'RE GOING ON A LEAF HUNT Leaf Book Video - abridged version*
*Phytomedicinal and Ameliorative Effect of Ethanol Leaf Extract of Corchorus olitorius on Carbon Leaf Book Video - Full Version*
*Phytochemical and Antiepileptic Activity of the Ethanol Leaf Extracts of Culsasia falcatifolia When the Leaf Blew In+ Fall Books for Kids+ Autumn Books Read Aloud*
My Leaf Book Insulin Plant ss Diabetes - Payo ni Doc Willie Ong #638 Story Time With Michele
1The Falling Leaves\* read aloud for kids
*Tea-Leaf-Reading—Dennis Fairehid—Witeheraft-Book Review*

Effects Of An Ethanolic Leaf
Effects of aqueous and ethanolic leaf extracts from drumstick tree (Moringa oleifera) on gilthead seabream (Sparus aurata L.) leucocytes, and their cytotoxic, antitumor, bactericidal and antioxidant activities

Effects of aqueous and ethanolic leaf extracts from ...
Abstract. Effects of ethanolic extracts of leaf, seed and fruit of Datura metel on kidney function of male albino rats was investigated in this study. The result showed a non-significant (p > 0.05) increase of urea concentration in groups 2 and 3 administered low (300 mg/kg bw) and high (600 mg/kg bw) dose of leaf extract respectively and a non-significant (p > 0.05) decrease in all groups administered the seed and fruit extracts compared with normal control (group 1).

Effects of ethanolic extracts of leaf, seed and fruit of ...
Leaf part is usually used to reduce the shortness of breath and as treatment of acne. The root part is used to treat headache and insomnia and is believed can accelerate fracture healing. Essential oil of J. sambacis used as fragrance for skin care products as it tones the skin as well as reduces skin inflammation.

Mechanisms of Gastroprotective Effects of Ethanolic Leaf ...
Mechanisms of Gastroprotective Effects of Ethanolic Leaf Extract of Jasminum sambac against HCl/Ethanol-Induced Gastric Mucosal Injury in Rats 1. Introduction. Peptic ulcer is a common disorder of the stomach and duodenum [ 1 D. A. O. V. Araujo, C. Takayama, F. M. 2. Materials and Methods. In this ...

Mechanisms of Gastroprotective Effects of Ethanolic Leaf ...
The 80% ethanolic leaf extracts of Musa paradisiaca (2.86 g/mL) and Musa acuminata (3.33 g/mL) inhibited the growth of MRSA and MSSA isolates but the extract of Musa sapientum (2.86 g/mL) had no effect against these bacteria.

Antibacterial effects of Musa sp. ethanolic leaf extracts ...
Abstract. The impact of Moringa oleifera leaf ethanol extract (MOLEE) was assessed on the expression of the steroidogenic genes (steroidogenic acute regulatory protein (StAR) and cytochrome P450c17 subfamily a (CYP17a) and luteinizing hormone receptor (LHR) gene) as well as on the cadmium chloride (CdCl2)-induced reproductive toxicity for 56 days in male rats.

Partial Ameliorative Effect of Moringa Leaf Ethanolic ...
Histology observation showed less edema and leucocytes infiltration as compared with the ulcer control which exhibited severe gastric mucosa injury. Furthermore, the leaf extract elevated the mucus weight, level of prostaglandin E2and superoxide dismutase. The extract also reduced malondialdehyde amount significantly.

Gastroprotective Effect of Ethanolic Extract of Curcuma ...
This study evaluated the effect of aqueous and ethanolic leaf extracts of Vitex doniana, against Streptozotocin (STZ) induced diabetic Wistar rats.Diabetes was induced with a single dose of Streptozotocin (60 mg kg-1 b.wt. i.p), followed by treatment with aqueous and ethanolic leaf extracts of V. doniana (100 and 200 mg kg-1 b.wt./day each), while metformin (25 mg kg-1 b.wt./day) was used as ...

Anthyperglycemic and Anthyperlipidemic Effects of ...
Several reports deduced that Moringa oleifera and Camellia sinensis have been shown to have antimicrobial effects against a variety of Gram positive and Gram negative bacteria (e.g., Escherichia coli, Salmonella spp., Staphylococcus aureus and Enterococcus spp.) and some fungi (e.g., Candida albicans) 7 - 9.

Antibacterial Effect of the Ethanol Leaves Extract of ...
Effects of ethanolic dried leaf extract of Lecaniodiscus cupanioides on antioxidant enzymes and biochemical parameters in rats. Oladimeji-Salami JA(1), Akindele AJ(2), Adeyemi OO(2). Author information: (1)Department of Pharmacology, Therapeutics and Toxicology, College of Medicine, University of Lagos, Idi-araba, Lagos, Nigeria.

Effects of ethanolic dried leaf extract of Lecaniodiscus ...
activity of an ethanolic leaf extract of P. guajava (guava) in vitro and in vivo. Our results demonstrated that guava leaf extract (GLE) significantly inhibited lipopolysaccharide (LPS)-induced production of nitric oxide and prostaglandin E2 in a dose-dependent manner. GLE suppressed the expression and activity of both

Anti-inflammatory effects of an ethanolic extract of guava ...
Effect of ethanolic leaf extract of noni (morinda citrifolia L) on blood pressure in ovariectomized rats fed with high-fat diet
Kumeshini Sukalingam, 1 Muhammad Zaid Zainuddin, 1 Gloria Chong, 1 Norhayati Aida, 2 Nurul Aina Abdullah, 2 Faizah Othman 1

Effect of ethanolic leaf extract of noni (morinda ...
Anticancer effects of ethanolic neem leaf extract on prostate cancer cell line (PC-3)
Prostate cancer (PC) is the most prevalent cancer and the leading cause of male cancer death. Azadirachta indica (neem tree) has been used successfully centuries to reduce tumors by herbalists throughout Southeast Asia. Here the present study indicated that an ethanolic extract of neem has been show ...

Anticancer effects of ethanolic neem leaf extract on ...
Subapiya and Nagini (2003) suggested that ethanolic neem leaf extract might exert its chromopreventive effects by modulating lipid peroxidation and enhancing the antioxidant status in the stomach, liver and erythrocytes.

Anticancer effects of ethanolic neem leaf extract on ...
To determine the effects of ethanolic leaf extract of Spondias mombin (S. mombin) on the histology of the anterior pituitary, ovary and uterus; and on the serum sex hormones of adult female Wistar rats.

Effects of ethanolic leaf extract of Spondias mombin on ...
The effect of ethanolic leaf extract of A. sclerocarpa (250 and 500 mg/kg, p.o.) was studied in experimentally induced renal stone in rats by in vivo model. Ethylene glycol model (0.75% in drinking water, for 28 days) was used for renal stone induction. The blood, urine and kidney samples were used for various parameters.

Protective effect of ethanolic leaf extract of Alphonsea ...
OBJECTIVE: The leaves of sage (Salvia officinalis L., Lamiaceae) are reported to have a wide range of biological activities, such as anti-bacterial, fungistatic, virustatic, astringent, eupeptic and anti-hydrotic effects. To determine the mnemogenic effect of sage leaves, we investigated the effects of ethanolic

Effects of Salvia officinalis L. (sage) leaves on memory ...
The results of the present study suggest that ethanolic O. sanctum leaf extract inhibits DMBA-induced genotoxicity and oxidative stress by modulating xenobiotic-metabolizing enzymes, reducing the extent of lipid and protein oxidation and up-regulating antioxidant defenses.

Ocimum sanctum Linn. (Holy Basil) ethanolic leaf extract ...
Histological effects of ethanolic leaf extract of Codiaeum variegatum on the cerebrum of adult Wistar rats
Elizabeth Finbarrs-Bello 1, Ogugua Augustine Egwu 2, Vivian Atuadu 1, Eni Ogbonnia Egwu 1 1 Department of Anatomy, College of Medicine, Ebonyi State University, Abakaliki, Ebonyi State, Nigeria 2 Department of Anatomy, Federal University Ndufu-Alike Ikwo, Ebonyi State, Nigeria

Arctocarpus ovatus Blanco is an endemic plant species belonging to the family Moraceae. Arctocarpus species have known medicinal value because they are rich in biological compounds such as flavonoids. This study evaluated the cholesterol-lowering activity of the A. ovatus ethanolic leaf extract (AOEE) in Sprague Dawley rats and its acute oral toxicity. The plant extract and its solvent fractions (test samples) were screened for the presence of secondary metabolites using qualitative tests and thin layer chromatography. The in vitro antioxidant activity of the test samples was determined using 2,2-diphenyl-1-picrylhydrazyl (DPPH), hydrogen peroxide and nitric oxide radical scavenging assays. The ethanolic leaf extract of A. ovatus was obtained through percolation using 95% ethanol as the extracting solvent and concentrated under reduced pressure. It was partitioned with hexane, dichloromethane (DCM) and n-butanol. The resulting fractions were also concentrated in vacuo. Phytochemical screening detected the presence of secondary metabolites such as anthraquinones, flavonoids, phenolics and terpenes in the ethanolic extract and DCM fraction. The AOEE and its solvent fractions were negative for alkaloids and saponins. In the in vitro antioxidant assays, the ethanolic extract demonstrated a significant DPPH (IC50 = 0.078 mg/mL) and nitric oxide (IC50 = 0.098 mg/mL) radical scavenging activities as well as hydrogen peroxide (IC50 = 0.045 mg/mL) scavenging effect. The ethanolic leaf extract of A. ovatus was subjected to acute oral toxicity test based on the guidelines of OECD 425 main test and was found to be safe and non-toxic up to 2000 mg/kg body weight of female Sprague Dawley rats. AOEE treated rats at doses of 200, 400 and 600 mg/kg showed time dependent reduction of serum levels of total cholesterol, triglycerides and low density lipoproteins after 14 days of oral administration (p

Bachelor Thesis from the year 2015 in the subject Chemistry - Bio-chemistry, grade: 3.84, course: Biochemistry, language: English, abstract: The present study was done to evaluate the acute (14 days) toxicity of the ethanolic leaf extract of Myrianthus arboreus on the liver enzymes of wistar rats. In the acute (14 days) toxicity studies, 24 rats were grouped into 1- 8 groups (n=3rats/cage) and administered with 1500, 1000 and 500 mg/kg body weight for 7 days and 14 days. The rats were sacrificed after 7 days and 14 day of administration and blood samples and liver organ were collected for investigations. The biochemical parameters such as the Alkaline phosphatase (ALP), Alanine transaminase (ALT) and Aspartate aminotransferase (AST) were determined and the liver histology analysed. The mean values of ALP showed significant increase (P=0.05) at groups 2, 3 and 4 and a significant increase (p=0.05) at all dosages and times except for group 2. The histological analysis showed microvesicular steatosis at groups 2 and 3 and a ballooning hepatic necrosis at group 7. The phytochemical analysis of Myrianthus arboreus shows the presence of alkaloids, flavonoids, tannins, anthraquinones, triterpenoids, carbohydrate, cardenolide and saponins in detectable limits but fixed oils and cyanogenic glycosides were not determined. In this investigation, we can conclude that the ethanolic leaf extract of Myrianthus arboreus was unsafe at all doses considered for a period of 14 days. However, at a dose below 500 mg for 7 days could be considered safe.

The leaves of Acanthopanax trifoliatus (L) Merr have been used to treat several diseases such as tuberculosis and lung hemorrhage, and as a tonic to improve general weakness. This study was conducted to determine the effect of ethanolic extract of Acanthopanax infoliatus. (L) Merr on both acute and chronic anti-inflammatory activity. The male Sprague-Dawley rats were randomly divided into five groups, with 6 animals per group for both acute and chronic inflammation. For inflammation study: (1) control group (5% ethanol); (2) group treated with 30 mg/kg of extract; (3) group treated with 100 mg/kg of extract; (4) group treated with 300 mg/kg of extract; (5) reference group: 30mg/kg of piroxicam for acute inflammation and 10 mg/kg of indomethacin for chronic inflammation. For acute inflammation study, the rats were injected subcutaneously with 0.1 ml of 1% carrageenan onto the plantar surface of right hind paw and equal volume of distilled water was injected onto the plantar surface of the left hind paw after 30 minute force fed of ethanolic extra of Acanthopanax trifolitus (L) Merr leaves (EAT).The volume of both hind of each rat was measured using a plethysmometer at every half- hourly interval until the period of five hours after the injection of the carrageenan. For chronic inflammation study, the rats were injected intradermally with 0.1ml of Freund's Complete Adjuvant (FCA) onto the right hind paw. The rats were acclimatized for 14 days after the adjuvant injection. The treatment was started on day 14 until day 28. The volume of the paw was measured before induction, before treatment, and after treatment using plethysmometer. The percentage of swelling and inhibition for all group in both acute and chronic inflammation was calculated and compared. For acute inflammation, EAT gave significant anti-inflammatory effect at EAT 300 mg/kg and piroxicam 30 mg/kg as compared to control group and the percentage of inhibition for the treated group of 30, 100 and 300 mg/kg were 10.50%, 32.43% and 46.23% respectively. For chronic inflammation, EAT exhibited significant effect also at 300 mg/kg and indomethacin 10 mg/kg as compared to control group and the percentage of inhibition for treatment group of 30, 100, 300 mg/kg were 9.57%, 23.71% and 47.57% respectively. It can be concluded that EAT exhibited a dose dependent inhibitory effect towards oedema in both acute and chronic infalammation. These result has shown that EAT leaves have anti-inflammatory properties. Further studies should be done to obtain effective dose (ED50) and may increase the dosage for EAT leaves treatment. Toxicity study (LD50) also should be carried out in order to determine the health safety usage in high dose concentration.

In vitro determination of the influence of salicylic acid on the anti-dandruff activity of an ethanolic extract of ketapeng leaves (Cassia alata L., Caesalpinaceae) was carried out according to the diffusion method using metal cylinders as reservoir. It was shown that this combination performed a synergistic effect against Staphylococcus aureus and Staphylococcus epidermidis at the minimum inhibition concentration of 0.75% salicylic acid and of 25% extract.A positive result could also be observed against fungi isolated from dandruff, as Mycosporum gypseum and Homodendron pedrosoi. [Authors' abstract].

Bachelor Thesis from the year 2015 in the subject Chemistry - Bio-chemistry, grade: 3.84, , course: Biochemistry, language: English, abstract: The present study was done to evaluate the acute (14 days) toxicity of the ethanolic leaf extract of Myrianthus arboreus on the liver enzymes of wistar rats. In the acute (14 days) toxicity studies, 24 rats were grouped into 1- 8 groups (n=3rats/cage) and administered with 1500, 1000 and 500 mg/kg body weight for 7 days and 14 days. The rats were sacrificed after 7 days and 14 day of administration and blood samples and liver organ were collected for investigations. The biochemical parameters such as the Alkaline phosphatase (ALP), Alanine transaminase (ALT) and Aspartate aminotransferase (AST) were determined and the liver histology analysed. The mean values of ALP showed significant increase (P=0.05), the ALT showed a non-significant increase (P=0.05) at groups 2, 3 and 4 and a significant increase (p=0.05) at groups 6, 7 and 8. The AST showed a non-significant increase (P=0.05) at all dosages and times except for group 2. The histological analysis showed microvesicular steatosis at groups 2 and 3 and a ballooning hepatic necrosis at group 7. The phytochemical analysis of Myrianthus arboreus shows the presence of alkaloids, flavonoids, tannins, anthraquinones, triterpenoids, carbohydrate, cardenolide and saponins in detectable limits but fixed oils and cyanogenic glycosides were not determined. In this investigation, we can conclude that the ethanolic leaf extract of Myrianthus arboreus was unsafe at all doses considered for a period of 14 days. However, at a dose below 500 mg for 7 days could be considered safe.

This book is focused on clarifying the anticancer effects (i.e., apoptotic, antiproliferative, antimetastatic, antiangiogenic) and mechanisms of most of the medicinal plants found in the world against solid and/or hematological cancers.

Unity in Diversity and the Standardisation of Clinical Pharmacy Services represents the proceedings of the 17th Asian Conference on Clinical Pharmacy (ACCP 2017), held 28—30 July 2017 in Yogyakarta, Indonesia. The primary aim of ACCP 2017 was to bring together experts from all fields of clinical pharmacy to facilitate the discussion and exchange of research ideas and results. The conference provided a forum for the dissemination of knowledge and exchange of experiences. As such, it brought together clinical pharmacy scholars, pharmacy practitioners, policy makers and stakeholders from all areas of pharmacy society and all regions of the world to share their research, knowledge, experiences, concepts, examples of good practice, and critical analysis with their international peers. This year also marks the celebration of 20 years of ACCP. Central themes of the conference and contributed papers were Clinical Pharmacy, Social and Administrative Pharmacy, Pharmacy Education, Pharmacoeconomics, Pharmacoepidemiology, Complementary and Alternative Medicine (CAM) and a number of related topics in the field of Pharmacy.

Copyright code : dbcc4049aac3b98584eb709e572049c5